55. Consider the following reaction :

$$X \xrightarrow{\text{(i) NaOH, CO}_2} \xrightarrow{\text{(ii) CHCl}_3 + \text{aq.NaOH}} Y$$

the products X and Y are

(A)
$$X = \bigcirc OH$$
 $COOH$ $Y = \bigcirc OH$

(B)
$$X = \bigcirc OH$$
 $COOH$

(C)
$$X = \bigcirc OH$$
 $Y = \bigcirc OH$ CHO

(D)
$$X = \bigcirc OH$$
 CHO

Case: Read the passage given below and answer the following questions (53-55).

Alcohols and Phenols are acidic in nature. Electron withdrawing groups in phenol increase its acidic strength and electron donating groups decrease it. Alcohols undergo nucleophilic substitution with hydrogen halides to give alkyl halides. On oxidation primary alcohols yield aldehydes with mild oxidising agents and carboxylic acids with strong oxidising agents while secondary alcohols yield ketones. The presence of –OH groups in phenols activates the ring towards electrophilic substitution. Various important products are obtained from phenol like salicylaldehyde, salicylic acid, picric acid etc.

53. Which of the following alcohols is resistant to oxidation?

- (B) CH₃–CH–OH CH₃
- (C) CH₃--CH₂--OH
- (D) CH₃-OH

54. Which of the following group increases the acidic character of phenol?

- (A) CH₃O-
- (B) CH₃-
- (C) NO_2-
- (D) All of these

Section - C

This section consists of 6 multiple choice questions with an overall choice to attempt any 5. In case more than desirable number of questions are attempted, ONLY first 5 will be considered for evaluation.

Match the following: 50.

latch the following:	
I	
(i) Stoichiometric defects	(a) Crystalline solids
(ii) long range order	(b) F-centres
(iii)ABC ABC ABC	(c) Schottky and Frenkel defects
(iv) Number of atoms per unit cell=2	(d) fcc structure
(v) Metal excess defect due to anionic vacancies	

Which of the following is the best matched options?

(A)
$$(i) - (d), (ii) - (a), (iii) - (b), (iv) - (c)$$

(B)
$$(i) - (c), (ii) - (a), (iii) - (d), (v) - (b)$$

(C)
$$(i) - (c), (ii) - (a), (iii) - (d), (iv) - (b)$$

(D)
$$(i) - (a), (ii) - (b), (v) - (c), (iv) - (d)$$

Which of the following analogies is correct? 51.

(A) XeF₂: linear :: XeF₆: square planar

(B) moist SO₂: Reducing agent :: Cl₂: bleaching agent

(C) N₂: Highly reactive gas :: F₂: inert at room temperature

(D) NH₃: strong base :: HI : weak acid

Complete the following analogy: 52.

Curdling of milk : A :: α-helix : B

B: Secondary structure (A) A: Primary structure

B: Primary structure (B) A: Denatured protein

B: Denatured protein (C) A: Secondary structure

B: Secondary structure (D) A: Denatured protein

- 47. Assertion (A): Chlorobenzene is less reactive towards nucleophilic substitution reaction.
 - **Reason** (R): Nitro group in chlorobenzene increases its reactivity towards nucleophilic substitution reaction.
 - (A) Both A and R are true and R is the correct explanation of A.
 - (B) Both A and R are true but R is not the correct explanation of A.
 - (C) A is true but R is false.
 - (D) A is false but R is true.
- 48. Assertion (A): Due to Schottky defect, there is no effect on the density of a solid.

 Reason (R): Equal number of cations and anions are missing from their normal sites in Schottky defect.
 - (A) Both A and R are true and R is the correct explanation of A.
 - (B) Both A and R are true but R is not the correct explanation of A.
 - (C) A is true but R is false.
 - (D) A is false but R is true.
- 49. Assertion (A): Fluorine forms only one oxoacid HOF.

Reason (R): Fluorine atom is highly electronegative.

- (A) Both A and R are true and R is the correct explanation of A.
- (B) Both A and R are true but R is not the correct explanation of A.
- (C) A is true but R is false.
- (D) A is false but R is true.

Which of the following isomer of pentane (C5H12) will give three isomeric monochlorides on photochemical chlorination?

- (B) CH₃ CH₂ CH₂ CH₂ CH₃
- (C) CH₃-CH-CH₂-CH₃
- (D) All of the above

Given below are the questions (45-49) labelled as Assertion (A) and Reason (R).

Select the most appropriate answer from the options given below:

Assertion (A): A raw mango placed in a saline solution loses water and shrivel into 45.

Reason (R): Through the process of reverse osmosis, raw mango shrivel into pickle.

- (A) Both A and R are true and R is the correct explanation of A.
- Both A and R are true but R is not the correct explanation of A. (B)
- A is true but R is false. (C)
- (D) A is false but R is true.

46.

Assertion (A): H₂S is less acidic than H₂Te.

Reason (R): H-S bond has more Δ_{bond} H° than H-Te bond

- (A) Both A and R are true and R is the correct explanation of A.
- Both A and R are true but R is not the correct explanation of A. (B)
- A is true but R is false. (C)
- A is false but R is true. (D)

Glucose on reaction with Br2 water gives: 10.

- (A) Saccharic acid
- (B) Hexanoic acid
- (C) Gluconic acid
- (D) Salicylic acid

Which of the following is optically inactive? 41.

- (A) (+) Butan-2-ol
- (B) (-) Butan-2-ol
- (C) (\pm) Butan–2–ol
- (D) (+) -2 Bromobutane


Which of the following is not a correct statement? 42.

- (A) Halogens are strong oxidising agents.
- (B) Halogens are more reactive than interhalogens.
- (C). All halogens are coloured.
- (D) Halogens have maximum negative electron gain enthalpy.

Which of the following has highest boiling point? 43.

- (A) C₂H₅-F
- (B) C₂H₅-Cl
- (C) C₂H₅-Br
- (D) C₂H₅-I

- Nucleic acids are polymer of
 - (A) amino acids
 - (B) nucleosides
 - (C) nucleotides
 - (D) glucose
- Which of the following gas dimerises to become stable? 37.
 - (A) CO₂ (g)
 - (B) NO₂ (g)
 - (C) SO₂ (g)
 - (D) N₂O (g)
- In the following diagram point, 'X' represents 38.



- Boiling point of solution
- Freezing point of solvent (B)
- Boiling point of solvent (C)
- (D) Freezing point of solution
- XeF₆ on reaction with NaF gives 39.
 - (A) $Na^{+}[XeF_{7}]^{-}$
 - $[NaF_2]^-[XeF_5]^+$ (B)
 - Na⁺[XeF₆]⁻ (C)
 - (D) $[NaF_2]^+ [XeF_5]^-$

Sulphuric acid is used to prepare more volatile acids from their corresponding salts due to its 31. due to its (A) strong acidic nature low volatility (B) (C) strong affinity for water (D) ability to act as a dehydrating agent An element with density 6 g cm⁻³ forms a fcc lattice with edge length of 4×10^{-8} cm. 32. The molar mass of the element is $(N_A = 6 \times 10^{23} \text{ mol}^{-1})$ (A) 57.6 g mol⁻¹ (B) 28.8 g mol⁻¹ (C) 82.6 g mol⁻¹ (D) 62 g mol⁻¹ In the reaction 33. compound 'Y' is OH -OMgBr

Which of the following is the weakest reducing agent in group 15?

The boiling point of a 0.2 m solution of a non-electrolyte in water is

(C)

(D)

(A) NH3 (B) PH₃ (C) AsH₃

(A) 100 °C

(C)

(B) 100.52 °C

(D) 100.26 °C

100.104 °C

(K_b for water = $0.52 \text{ K/kg mol}^{-1}$)

34.

35.

section consists of 24 multiple choice questions with overall choice to attempt any 20 mill be strong. In case more than desirable multiple choice questions with overall choice to attempt any 20 mill be This solutions. In case more than desirable number of questions are attempted, ONLY first 20 will be oldered for evaluation. gunsidered for evaluation.

- An unknown gas 'X' is dissolved in water at 2.5 bar pressure and has mole fraction 0.04 in solution. The mole fraction of 'X' gas when the pressure of gas is doubled at the same temperature is
 - (A) 0.08
 - (B) 0.04
 - (C) 0.02
 - (D) 0.92
- The base which is present in DNA but not in RNA, is
 - (A) Cytosine
 - (B) Guanine
 - (C) Adenine
 - (D) Thymine
- In the following reaction 28.

 CH_3 -CH=CH- CH_2 -OH \xrightarrow{PCC} the product formed is

- (A) CH3-CHO and CH3CH2OH
- (B) CH₃-CH=CH-COOH
- (C) CH₃-CH=CH-CHO
- (D) CH₃-CH₂-CH₂-CHO
- Enantiomers differ only in 29.
 - (A) boiling point
 - (B) rotation of polarised light
 - (C) melting point
 - (D) solubility
- The number of lone pairs of electrons in XeF4 is 30.
 - (A) zero
 - (B) one
 - (C) two
 - (D) three

Which of the following is a network solid?

(A) SO2

SiO2 (B)

(C) CO2

(D) · H2O

Major product formed in the following reaction

CH₃

$$CH_3 - C - Br + NaOCH_3 \rightarrow CH_3$$

$$CH_3$$

(D)
$$CH_3-C=CH_2$$

- Chlorine reacts with cold and dilute NaOH to give 24.
 - (A) NaCl and NaClO₃
 - (B) NaCl and NaClO
 - (C) NaCl and NaClO4
 - (D) NaClO and NaClO3
- Elevation of boiling point is inversely proportional to 25.
 - (A) molal elevation constant (Kb)
 - (B) molality (m)
 - (C) molar mass of solute (M)
 - (D) weight of solute (W)

Consider the following compounds: 17.

$$CI$$
, CH_{2} - CI , CI

the correct order of reactivity towards S_{N2} reaction (A) 1 > 111 > 11

- (B) 11 > 111 > 1
- (C) 11 > 1 > 111
- (D) 111 > 1 > 11

Which of the following forms strong pπ-pπ bonding? 18.

- (A) S8
 - (B) Ses
 - (C) Tes
 - (D) O2

19. F2 acts as a strong oxidising agent due to

- (A) low Δbond H° and low Δhyd H°
- (B) low Δbond H° and high Δhyd H°
- (C) high Δbond H° and high Δeg H°
- (D) low Δ_{hyd} H° and low Δ_{eg} H°

20. Which of the following sugar is known as dextrose?

- (A) Glucose
- (B) Fructose
- (C) Ribose
- (D) Sucrose

21. Cu reacts with dilute HNO3 to evolve which gas?

- (A) N₂O
- (B) NO₂
- (C) NO
- (D) N₂

	Which of the following acids reacts with acetic anhydride to form a compound Aspirin? (A) Benzoic acid (B) Salicylic acid (C) Phthalic acid (D) Acetic acid
13.	 Which of the following statements is wrong? (A) Oxygen shows pπ – pπ bonding. (B) Sulphur shows little tendency of catenation. (C) Oxygen is diatomic whereas sulphur is polyatomic. (D) O-O bond is stronger than S-S bond.
14.	diet are known as (A) Acidic amino acids (B) Essential amino acids (C) Basic amino acids (D) Non-essential amino acids
15.	Which one of the following halides contains C _{sp} ² -X bond? (A) Allyl halide (B) Alkyl halide (C) Benzyl halide (D) Vinyl halide (D) Vinyl halide

- Nucleosides are composed of (A) a pentose sugar and phosphoric acid 6. (B) a nitrogenous base and phosphoric acid (C) a nitrogenous base and a pentose sugar (D) a nitrogenous base, a pentose sugar and phosphoric acid The oxidation state of -2 is most stable in : (A) O (B) S (C) Se (D) Te Which of the following is not a characteristic of a crystalline solid? (A) A true solid (B) A regular arrangement of constituent particles (C) Sharp melting point (D) Isotropic in nature Which of the following formula represents Raoult's law for a solution containing non-volatile solute? (A) $p_{solute} = p^{\circ}_{solute}$. Xsolute (B) $p = K_{H,X}$ (C) $P_{Total} = p_{solvent}$ (D) $p_{solute} = p^{\circ}_{solvent}$. Xsolvent 10. An azeotropic solution of two liquids has a boiling point lower than either of the two when it (A) shows a positive deviation from Raoult's law. (B) shows a negative deviation from Raoult's law.
 - (C) shows no deviation from Raoult's law.
 - (D) is saturated.
- 11. Which of the following crystal will show metal excess defect due to extra cation?
 - (A) AgCl
 - (B) NaCl
 - (C) FeO
 - (D) ZnO

The structure of pyrosulphuric acid is O

$$(D) \quad HO - S - O - O - S - OH$$

The C-O-H bond angle in alcohol is

- (A) slightly greater than 109°28'.
- (B) slightly less than 109°28'.
- (C) slightly greater than 120°.
- (D) slightly less than 120°.

5. Consider the following reaction:

CH₃-CH=CH₂
$$\frac{1. \text{HBr}}{2. \text{ aq. KOH}}$$

The major end product is

General Instructions : This question paper contains 55 questions out of which 45 questions are to be attempted

- All questions carry equal marks. 1.
- This question paper consists of three Sections Section A, B and C. 2.
- Section A contains 25 questions. Attempt any 20 questions from Q. No. 01 to 25 3.
- Section B contains 24 questions. Attempt any 20 questions from Q. No. 26 to 49 4.
- Section C contains 6 questions. Attempt any 5 questions from Q. No. 50 to 55. 5.
- The first 20 questions attempted in Section A and Section B and first 5 questions 6. attempted in Section C by a candidate will be evaluated.
- There is only one correct option for every multiple choice question (MCQ). Marks will not be awarded for answering more than one option.
- There is no negative marking.

Section - A

This section consists of 25 multiple choice questions with overall choice to attempt any 20 questions. In case more than desirable number of questions are attempted, ONLY first 20 will be considered for evaluation.

- Which one of the following pairs will form an ideal solution?
- (A) Chloroform and acetone
- (B) Ethanol and acetone
- (C) n-hexane and n-heptane
- (D) Phenol and aniline
- Which of the following is known as amorphous solid?
- (A) Glass
- (B) Plastic
- (C) Rubber
- (D) All of the above

[P.T.O.