



Trigonal bipyramidal

MI₅



For example
(refer to VSEPR Theory !)




Example: $\text{Ru}(\text{CO})_3(\text{PPh}_3)_2$

Which possible structures can we draw from this compound and which one is most likely to form ?



Popular example: Iron Pentacarbonyl

Organic reactions on Cyclobutadiene:



Compare d^{10} and d^5

Both kinds of electron configuration result in zero CFSE.

What is the difference ?

Draw the energy level diagram for both answers and state if high- or lowspin

Which of the following compounds has a CFSE of $0.0\Delta_o$ associated with it?

- A. $[\text{PtF}_6]$
- B. $[\text{Fe}(\text{CN})_6]^{3-}$
- C. $[\text{Mn}(\text{OH}_2)_6]^{2+}$


Which of the following compounds has a CFSE of $-1.2\Delta_t$ associated with it?

- A. $[\text{CoCl}_4]^{2-}$
- B. $[\text{ReO}_4]^-$
- C. $[\text{FeCl}_4]^-$

UV/VIS Spectroscopy


Light absorption by electron transfers

Visible Colours



Colour of light <i>absorbed</i>	Approximate wavelength ranges / nm	Colour of light <i>transmitted</i>
Red	700-620	Green
Orange	620-580	Blue
Yellow	580-560	Violet
Green	560-490	Red
Blue	490-430	Orange
Violet	430-380	Yellow

UV-visible spectrum of $[\text{Ti}(\text{OH}_2)_6]^{3+}$



How to convert between cm^{-1} and nm ?

Why do we use wavenumber ν instead of wavelength λ ?

Question

- Which colour will the **S** complex have ?
- How many electron transitions are happening and why ?
- How will the colour change if we add ammonia to the aqueous solution ? And why ?

Compare 2 Ni-compounds


Ni(H₂O)₆ (2+)

Colour:


Ni(NH₃)₆ (2+):

Colour:

Which compound absorbs light at higher energy ?




Jahn-Teller Effect



The Jahn-Teller effect is generally only important for odd number occupancy of the e_g level.

The effect of Jahn-Teller distortions is best documented for Cu(II) complexes (with 3 electrons in the e_g level) where the result is that most complexes are found to have elongation along the z-axis.



Complexes



- A central metal atom bonded to a group of molecules or ions is a **metal complex**.
- If it's charged, it's a **complex ion**.
- Compounds containing complexes are **coordination compounds**.

Complexes

- The molecules or ions coordinating to the metal are the **ligands**.
- They are usually anions or polar molecules.
- They must have lone pairs to interact with metal



A chemical mystery:


Same metal, same

TABLE 24.1 Properties of Some Ammonia Complexes of Cobalt(III)

Original Formulation	Color	Ions per Formula Unit	"Free" Cl^- Ions per Formula Unit	Modern Formulation
$\text{CoCl}_3 \cdot 6 \text{NH}_3$	Orange	4	3	$[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
$\text{CoCl}_3 \cdot 5 \text{NH}_3$	Purple	3	2	$[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Green	2	1	<i>trans</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Violet	2	1	<i>cis</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$

- Many coordination compounds are brightly colored, but again, same metal, same ligands, different colors.

Werner's Theory



Co(III) oxidation state

Coordination # is 6

- suggested in 1893 that metal ions have **primary** and **secondary** valences.
 - Primary valence equal the metal's oxidation number
 - Secondary valence is the number of atoms directly bonded to the metal (coordination number)

Werner's Theory

- The central metal and the ligands directly bonded to it make up the **coordination sphere** of the complex.
- In $\text{CoCl}_3 \cdot 6 \text{NH}_3$, all six of the ligands are NH_3 and the 3 chloride ions are outside the coordination sphere.

TABLE 24.1 Properties of Some Ammonia Complexes of Cobalt(III)

Original Formulation	Color	Ions per Formula Unit	"Free" Cl^- Ions per Formula Unit	Modern Formulation
$\text{CoCl}_3 \cdot 6 \text{NH}_3$	Orange	4	3	$[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
$\text{CoCl}_3 \cdot 5 \text{NH}_3$	Purple	3	2	$[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Green	2	1	<i>trans</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Violet	2	1	<i>cis</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$

Chemistry of
Coordination
Compounds

Werner's Theory

In $\text{CoCl}_3 \cdot 5 \text{NH}_3$ the five NH_3 groups and one chlorine are bonded to the cobalt, and the other two chloride ions are outside the sphere.

TABLE 24.1 Properties of Some Ammonia Complexes of Cobalt(III)

Original Formulation	Color	Ions per Formula Unit	"Free" Cl^- Ions per Formula Unit	Modern Formulation
$\text{CoCl}_3 \cdot 6 \text{NH}_3$	Orange	4	3	$[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
$\text{CoCl}_3 \cdot 5 \text{NH}_3$	Purple	3	2	$[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Green	2	1	<i>trans</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Violet	2	1	<i>cis</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$

Chemistry of
Coordination
Compounds

Werner's Theory



Werner proposed putting all molecules and ions within the sphere in brackets and those “free” anions (that dissociate from the complex ion when dissolved in water) outside the brackets.

TABLE 24.1 Properties of Some Ammonia Complexes of Cobalt(III)

Original Formulation	Color	Ions per Formula Unit	“Free” Cl^- Ions per Formula Unit	Modern Formulation
$\text{CoCl}_3 \cdot 6 \text{NH}_3$	Orange	4	3	$[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
$\text{CoCl}_3 \cdot 5 \text{NH}_3$	Purple	3	2	$[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Green	2	1	<i>trans</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
$\text{CoCl}_3 \cdot 4 \text{NH}_3$	Violet	2	1	<i>cis</i> - $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$

Chemistry of
Coordination
Compounds


Werner's Theory



- This approach correctly predicts there would be two forms of $\text{CoCl}_3 \cdot 4 \text{NH}_3$.
 - The formula would be written $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$.
 - One of the two forms has the two chlorines next to each other.
 - The other has the chlorines opposite each other.

What is Coordination?

- When an orbital from a ligand with lone pairs in it overlaps with an empty orbital from a metal




Sometimes called a
coordinate covalent
bond

So ligands *must* have lone pairs of electrons.

Metal-Ligand Bond

- This bond is formed between a Lewis acid and a Lewis base.
 - The ligands (Lewis bases) have nonbonding electrons.
 - The metal (Lewis acid) has empty orbitals.




Metal-Ligand Bond

The metal's coordination ligands and geometry can greatly alter its properties, such as color, or ease of oxidation.



Chemistry of
Coordination
Compounds

Oxidation Numbers




Knowing the charge on a complex ion and the charge on each ligand, one can determine the oxidation number for the metal.

Oxidation Numbers





Or, knowing the oxidation number on the metal and the charges on the ligands, one can calculate the charge on the complex ion.

Example: $\text{Cr(III)(H}_2\text{O)}_4\text{Cl}_2$

$$+3 + 4(0) + 2(-1) = +1$$







Coordination Number



- The atom that supplies the lone pairs of electrons for the metal-ligand bond is the **donor atom**.
- The number of these atoms is the **coordination number**.

Coordination Number




- Some metals, such as chromium(III) and cobalt(III), consistently have the same coordination number (6 in the case of these two metals).
- The most commonly encountered numbers are 4 and 6.


Geometries

- There are two common geometries for metals with a coordination number of four:

- Tetrahedral
- Square planar





Tetrahedral



Square planar


Why square planar? We'll get to that

Geometries



By far the most-encountered geometry, when the coordination number is six, is octahedral.


Polydentate Ligands



- Some ligands have two or more donor atoms.
- These are called **polydentate ligands** or **chelating agents**.
- In ethylenediamine, $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$, represented here as en, each N is a donor atom.
- Therefore, en is **bidentate**.


Polydentate Ligands

Ethylenediaminetetraacetate, mercifully abbreviated EDTA, has six donor atoms.











CoEDTA⁻

Wraps around the central atom like an octopus




[EDTA]⁴⁻

Polydentate Ligands

Ligand Type	Examples
Monodentate	$\text{H}_2\ddot{\text{O}}:$ Water $:\ddot{\text{F}}:^-$ Fluoride ion $:\text{C}\equiv\text{N}:^-$ Cyanide ion $:\ddot{\text{O}}-\text{H}^-$ Hydroxide ion $:\text{NH}_3$ Ammonia $:\ddot{\text{Cl}}:^-$ Chloride ion $[\text{:}\ddot{\text{S}}=\text{C}=\ddot{\text{N}}:]^-$ Thiocyanate ion $[\text{:}\ddot{\text{O}}=\text{N}=\ddot{\text{O}}:]^-$ Nitrite ion
Bidentate	 Ethylenediamine (en)  Bipyridine (bipy)  <i>Ortho</i> -phenanthroline (<i>o</i> -phen)  Oxalate ion  Carbonate ion
Polydentate	 Diethylenetriamine  Triphosphate ion  Ethylenediaminetetraacetate ion (EDTA^{4-})

Chelating agents generally form more stable complexes than do monodentate ligands.


Chelating Agents




- Bind to metal ions removing them from solution.
- Phosphates are used to tie up Ca^{2+} and Mg^{2+} in hard water to prevent them from interfering with detergents.

Chelating Agents

- Porphyrins are complexes containing a form of the porphine molecule shown at right.
- Important biomolecules like heme and chlorophyll are porphyrins.



Chelating Agents



Porphines (like chlorophyll *a*) are tetradentate ligands.


Nomenclature of

Ligand	Name in Complexes	Ligand	Name in Complexes
Azide, N_3^-	Azido	Oxalate, $\text{C}_2\text{O}_4^{2-}$	Oxalato
Bromide, Br^-	Bromo	Oxide, O^{2-}	Oxo
Chloride, Cl^-	Chloro	Ammonia, NH_3	Ammine
Cyanide, CN^-	Cyano	Carbon monoxide, CO	Carbonyl
Fluoride, F^-	Fluoro	Ethylenediamine, en	Ethylenediamine
Hydroxide, OH^-	Hydroxo	Pyridine, $\text{C}_5\text{H}_5\text{N}$	Pyridine
Carbonate, CO_3^{2-}	Carbonato	Water, H_2O	Aqua

- The basic protocol in coordination nomenclature is to name the ligands attached to the metal as prefixes before the metal name.
- Some common ligands and their names are listed above.


Nomenclature of Coordination Compounds

- As always the name of the **cation** appears first; the **anion** is named last.
- Ligands are listed alphabetically before the metal. Prefixes denoting the number of a particular ligand are ignored when alphabetizing.




Nomenclature of Coordination Compounds

- The names of anionic ligands end in “o”; the endings of the names of neutral ligands are not changed.
- Prefixes tell the number of a type of ligand in the complex. If the name of the ligand itself has such a prefix, alternatives like *bis-*, *tris-*, etc., are used.






Nomenclature of Coordination Compounds

- If the complex is an anion, its ending is changed to *-ate*.
- The oxidation number of the metal is listed as a Roman numeral in parentheses immediately after the name of the metal.



Examples of complexes derived





from



Ligand	Complex	IUPAC name
Water	$[\text{Cr}(\text{OH}_2)_6]^{3+}$	Hexaaquachromium(III)
Halide	$\text{K}[\text{FeCl}_4]$	Potassium tetrachloroferrate
Ammonia	$[\text{Ag}(\text{NH}_3)_2]^+$	Diamminesilver(I)
Cyanide	$[\text{Ni}(\text{CN})_4]^{2-}$	Tetracyanonickelate(II)


Examples of complexes derived

from



Ligand	Complex	IUPAC name
pyridine	$\text{Ru}(\text{py})_4\text{Cl}_2$	Tetrapyridineruthenium(II) dichloride
Ethylene diamine	$[\text{Co}(\text{en})_3]^{3+}$	Tris(1,2-diaminoethane)cobalt(III)
Bipyridine (bipy)	$[\text{Ru}(\text{bipy})_3]^{3+}$	Tris(2,2'-bipyridyl)ruthenium(III)
oxine	-	8-hydroxyquinolinato


Isomers




Isomers have the same molecular formula, but their atoms are arranged either in a different order (structural isomers) or spatial arrangement (stereoisomers).

Structural Isomers

If a ligand (like the NO_2 group at the bottom of the complex) can bind to the metal with one or another atom as the donor atom, linkage isomers are formed.



Nitro isomer




Nitrito isomer

Chemistry of
Coordination
Compounds

Structural Isomers

- Some isomers differ in what ligands are bonded to the metal and what is outside the coordination sphere; these are **coordination-sphere isomers**.
- Three isomers of $\text{CrCl}_3(\text{H}_2\text{O})_6$ are
 - The violet $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3$,
 - The green $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2 \cdot \text{H}_2\text{O}$, and
 - The (also) green $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl} \cdot 2 \text{H}_2\text{O}$.

Geometric isomers



- With these geometric isomers, two chlorines and two NH_3 groups are bonded to the platinum metal, but are clearly different.



- *cis*-Isomers have like groups on the same side.
- *trans*-Isomers have like groups on opposite sides.

of each atom the same

Bonding the same

Arrangement in space different


Stereoisomers



- Other stereoisomers, called **optical isomers** or **enantiomers**, are mirror images of each other.
- Just as a right hand will not fit into a left glove, two enantiomers cannot be superimposed on each other.


Enantiomers

A molecule or ion that exists as a pair of enantiomers is said to be **chiral**.




Enantiomers



- Most of the physical properties of chiral molecules are the same, boiling point, freezing point, density, etc.
- One exception is the interaction of a chiral molecule with plane-polarized light.





Enantiomers

- If one enantiomer of a chiral compound is placed in a polarimeter and polarized light is shone through it, the plane of polarization of the light will rotate.
- If one enantiomer rotates the light 32° to the right, the other will rotate it 32° to the left.






Copyright © 2006 Pearson Prentice Hall, Inc.



Copyright © 2006 Pearson Prentice Hall, Inc.

Chemistry of
Coordination
Compounds



Copyright © 2006 Pearson Prentice Hall, Inc.

Chemistry of
Coordination
Compounds

Explaining the properties of transition metal coordination complexes

1. Magnetism

2. color

Metal complexes and color

The ligands of a metal complex effect its color




Copyright © 2006 Pearson Prentice Hall, Inc.

Addition of NH_3 ligand to $\text{Cu}(\text{H}_2\text{O})_4$ changes its color

Chemistry of
transition
metals

Why does anything have color?



Copyright © 2006 Pearson Prentice Hall, Inc.

Light of different frequencies give different colors

We learned that elements can *emit* light of different frequency or color.

But these coordination complexes are not emitting light

They *absorb* light.

How does that give color?

Chemistry of
Coordination
Compounds

Light can bounce off an object or get absorbed by object

No light absorbed, all reflected get white color
All light absorbed, none reflected get Black color
What if only one color is absorbed?

Complimentary color wheel


If one color absorbed, the color opposite is perceived.

Absorb **Orange**

See **Blue**

Absorb **Red**

See **Green**




Copyright © 2006 Pearson Prentice Hall, Inc.



Absorbs in
green yellow.
Looks purple.




(a)



(b)

A precise measurement of the absorption spectrum of Compounds is critical



Copyright © 2006 Pearson Prentice Hall, Inc.

Metal complexes and color

But why do different ligands on same metal give Different colors?

Why do different ligands change absorption?




Copyright © 2006 Pearson Education, Inc.

Addition of NH_3 ligand to $\text{Cu}(\text{H}_2\text{O})_4$ changes its color

Chemistry of
transition
metals

Model of ligand/metal bonding. Electron pair comes from ligand Bond very polarized.

Assumption: interaction pure electrostatic.



Copyright © 2006 Pearson Prentice Hall, Inc.